

SEPHU triage risk stratification assessment

	Low Risk	Medium (clinically well)	Medium (clinically unwell)	High
	Need age, symptom and social to be met	Only one of Age, Comorbidity or Unwell symptoms need to be met		
Age	<60	>60	Any	Any
	Asymptomatic or mild symptoms (refer to questionnaire script)	Asymptomatic or mild symptoms	New dyspnoea Persistent cough Significant systemic symptoms – ie fever>40C, sweats, chills, repeated diarrhoea, nausea, vomiting, dehydration	SOB at rest Chest Pain Syncope Altered mental state Haemoptysis Clinical Concern
Comorbidities	Controlled Hypertension Well controlled diabetes Obesity Active smoker <15cpd Ex smoker Alcohol Dependency# Substance Abuse# Mental health #	Poorly controlled Hypertension Cardiovascular disease Respiratory disease Poorly controlled diabetes Immunocompromised (chemotherapy, steroids, immunosuppressants) Malignancy CKD CLD Active smoker >15cpd Neurological Disease Haematological disease Pregnancy* Alcohol Dependency and or Substance abuse# Mental health #		Any
Home isolation	Barriers to home isolation will be reviewed and appropriate support mechanisms put in place			
Patient consent	Patient consents to Referral through pathway			

* Pregnant patients will need to be escalated to Monash Health's pathway

These patients can be managed in the community. Further assessments and support are available via Star Health

NOTES:

- If clinical concern, GP to be alerted and extra assessment to be completed, and escalation of pathway as appropriate
- Clearance team as per SEPHU/DoH guidelines and protocols
- ATSI consider age <50 or >50
- Clinician discretion required in stratification
- Barrier to home isolation may be cause for Alfred Medium risk stream